



DEAS 768:2016

DRAFT EAST AFRICAN STANDARD

Fortified milled maize (corn) products — Specification

EAST AFRICAN COMMUNITY

Copyright notice

This EAC document is copyright-protected by EAC. While the reproduction of this document by participants in the EAC standards development process is permitted without prior permission from EAC, neither this document nor any extract from it may be reproduced, stored or transmitted in any form for any other purpose without prior written permission from EAC.

Requests for permission to reproduce this document for the purpose of selling it should be addressed as shown below or to EAC's member body in the country of the requester:

*© East African Community 2013 — All rights reserved
East African Community
P.O. Box 1096
Arusha
Tanzania
Tel: 255 27 2504253/8
Fax: 255 27 2504481/2504255
E-mail: eac@eachq.org
Web: www.eac-quality.net*

Reproduction for sales purposes may be subject to royalty payments or a licensing agreement.
Violators may be persecuted

Table of contents

Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references	1
3 Terms and definitions	1
4 Quality requirements.....	2
5 Fortification requirements.....	3
6 Food additives.....	5
7 Hygiene.....	5
8 Contaminants	5
9 Packaging.....	6
10 Labelling.....	6
11 Methods of sampling.....	7
12 Methods of Testing	7
Annex A (normative) Determination of acid insoluble ash.....	9
Bibliography	10

Foreword

Development of the Draft East African Standard has been necessitated by the need for harmonizing requirements governing quality of products and services in the East African Community. It is envisaged that through harmonized standardization, trade barriers which are encountered when goods and services are exchanged within the Community will be removed.

In order to meet the above objectives, the EAC Partner States have enacted an Draft East African Standardization, Quality Assurance, Metrology and Testing Act, 2006 (EAC SQMT Act, 2006) to make provisions for ensuring standardization, quality assurance, metrology and testing of products produced or originating in a third country and traded in the Community in order to facilitate industrial development and trade as well as helping to protect the health and safety of society and the environment in the Community.

Draft East African Standards are formulated in accordance with the procedures established by the Draft East African Standards Committee. The Draft East African Standards Committee is established under the provisions of Article 4 of the EAC SQMT Act, 2006. The Committee is composed of representatives of the National Standards Bodies in Partner States, together with the representatives from the private sectors and consumer organizations. Draft Draft East African Standards are circulated to stakeholders through the National Standards Bodies in the Partner States. The comments received are discussed and incorporated before finalization of standards, in accordance with the procedures of the Community.

Article 15(1) of the EAC SQMT Act, 2006 provides that “Within six months of the declaration of an Draft East African Standard, the Partner States shall adopt, without deviation from the approved text of the standard, the Draft East African Standard as a national standard and withdraw any existing national standard with similar scope and purpose”.

Draft East African Standards are subject to review, to keep pace with technological advances. Users of the Draft East African Standards are therefore expected to ensure that they always have the latest versions of the standards they are implementing.

© East African Community 2013 — All rights reserved*
East African Community
P O Box1096
Arusha
Tanzania
Tel: 255 27 2504253/8
Fax: 255-27-2504481/2504255
E-Mail: eac@eachq.org
Web: www.each.org

Introduction

The Health Ministers of the East, Central and Southern Africa (ECSA) Health Community passed a resolution in 2002 directing the Secretariat to work with the countries to fortify commonly consumed foods in the region after recognizing the high levels of malnutrition in the region. ECSA-HC is an inter-governmental organization that fosters cooperation in health among countries in the East, Central and Southern African Region. It has 10 active member states namely Kenya, Uganda, Tanzania, Malawi, Zambia, Zimbabwe, Lesotho, Swaziland, Mauritius and Seychelles. The mandate of the organization is to promote relevance and efficiency in health in the region.

Following initial promotion efforts, the countries identified staple foods suitable for fortification as oil, sugar, maize meal/ flour and wheat flour. These foods can be used as vehicles to deliver essential micronutrients to the populations. Based on scientific evidence and working with countries using country data, the Secretariat developed implementation focused guidelines on fortification of these foods to help countries start up programs and scale up the existing programs. These guidelines included fortification levels for addition of micronutrients at the factory, and levels for monitoring at commercial level.

Based on the guidelines and other available information, most of the countries in the East African Region and in the larger Africa have initiated national programs on oil fortification with vitamin A; and wheat and maize meal/ flour fortification with iron, zinc, folic acid, niacin, vitamin B-1, B-2, B6 and B-12 and vitamin A. Sugar fortification with vitamin A has also been considered as a way of supplementing other sources of the vitamin in order to prevent and reduce problems associated with the deficiency of this vitamin. Salt fortification with iodine continues to be implemented in all the countries.

With the increased trade of food commodities including these fortified foods within the region, it has become imperative to develop regional standards that over and above the other standards, stipulate minimum and maximum levels of the added nutrients, provide clauses on how to pack the fortified product and the use of health and nutrition claims. The guidelines developed through ECSA have now been incorporated into food standards to provide for specific fortified products.

It is envisaged that, the adoption of these standards and their utilization within the region will help countries adopt food fortification as a strategy to prevent, alleviate or eliminate micronutrient deficiency in the region. Standards will not only promote the health of the population but will also ensure safety of food products and enhance fair trade.

This standard was developed with support from the East, Central and Southern Africa Health Community (ECSA-HC) Secretariat. This was possible through a grant by the A2Z Project of the United States Agency for International Development (USAID). The financial and technical support was used in the process of formulation of fortification levels, development of the draft standards and mobilization of stakeholders to review the standard in national and regional fora. This support is hereby acknowledged.

Fortified milled maize (corn) products — Specification

1 Scope

This Draft East African Standard specifies the requirements and methods of sampling and test for fortified milled maize (corn) products namely whole maize meal, granulated maize meal, sifted maize meal and maize flour from the grains of common maize (*Zea mays L.*) intended for human consumption.

2 Normative references

The following normative documents contain provisions which, through reference in this text constitute provisions of this Draft East African Standard;

EAS 2, *Maize (grains) — Specification*

EAS 38, *Labelling of pre-packaged foods — Specification*

EAS 39, *Code of practice for hygiene in the food and drink manufacturing industry*

EAS 44, *Milled maize products — Specification*

EAS 103, *General standard for food additives*

CAC/GL 1, *General guidelines on claims*

CAC/GL 2, *Guidelines on nutrition labelling*

CAC/GL 23, *Guidelines for use of nutrition and health claims*

CODEXSTAN 193, *Codex general standards for contaminants and toxins in food and feed*

ISO 711, *Cereals and cereal products — Determination of moisture content (Basic reference method)*

ISO 6540, *Maize — Determination of moisture content (on milled grains and on whole grains)*

ISO 5498, *Agricultural food products — Determination crude fibre content-General method*

ISO 6579, *Microbiology of food and animal feeding stuffs — Horizontal method for the detection of *Salmonella* spp.*

ISO 7251, *Microbiology of food and animal feeding stuffs — Horizontal method for the detection and enumeration of presumptive *Escherichia coli* — Most probable number technique*

ISO 7305, *Milled cereal products – Determination of fat acidity*

ISO 11085, *Cereals, cereals-based products and animal feeding stuffs -- Determination of crude fat and total fat content by the Randall extraction method*

ISO 13690, *Cereals, pulses and milled products — Sampling of static batches*

ISO 16050, *Foodstuffs — Determination of aflatoxin B1, and the total content of aflatoxins B1, B2, G1 and G2 in cereals, nuts and derived products — High-performance liquid chromatographic method*

ISO 20483, *Cereals and pulses — Determination of the nitrogen content and calculation of the crude protein content — Kjeldahl method*

ISO 21527-2, *Microbiology of food and animal feedstuffs — Horizontal method for the enumeration of yeasts and moulds — Part 2: Colony count technique in products with water activity less than or equal to 0.95*

3 Terms and definitions

For the purposes of this standard the definitions in EAS 44 and the following shall apply.

3.1

diluent

suitable, inert, edible food-grade carrier for micronutrients

3.2

premix

blend of fortificants and diluents formulated to provide specified and determinable amounts of micronutrients

3.3

fortified milled maize products

maize meal, maize flour or sifted maize meal to which micronutrients have been added in accordance with this standard

3.4

fortificant

compound which contains the specified micronutrient intended to be added to a food vehicle

3.5

food fortification

practice of deliberately adding essential micronutrients in a food to improve the nutritional quality of the food and to provide a public health benefit with minimal risk to health

3.6

food grade container

container which will safeguard the hygienic, nutritional, technological, and organoleptic qualities of the product

4 Requirements

4.1 General requirements

4.1.1 Maize grain

Fortified milled maize (corn) products shall be made from maize grain complying with EAS 2.

4.1.2 Fortified milled maize (corn) products shall;

- a) be of characteristic colour conforming to the colour of maize from which it was prepared;
- b) not contain any foreign matter such as insects, fungi, dirt or other contaminants;
- c) be free from fermented musty or other objectionable odour and colours;
- d) be wholesome and fit for human consumption in all aspects.

4.2 Specific requirements

Fortified milled maize (corn) products shall conform to the requirements given in Table 1.

Table 1 — Specific quality requirements for fortified milled maize (corn) products

SI No.	Characteristic	Type of product				
		Sifted maize meal	Granulated maize meal	Whole maize meal	Maize flour	Test methods
i)	Crude fibre content, % by mass, max.	0.7	1.0	3.0	0.7	ISO 5498
ii)	Crude fat on a moisture free basis, % by mass, max.	3.0	3.0	3.1*	3.0	ISO 11085
iii)	Moisture content, % by mass, max.	14	14	14	14	ISO 6540
iv)	Total ash, % by mass, max.	1.0	1.0	3.0	1.0	ISO 2171
v)	Acid insoluble ash, % by mass, max.	0.15	0.35	0.40	0.15	Annex A
vi)	Fat acidity, mg KOH per 100g of product, on dry matter basis, max.	80	80	80	80	ISO 7305
*Minimum crude fat on moisture free basis for whole maize meal						

5 Fortification requirements

5.1 Levels of micronutrients

The fortified milled maize products shall conform to the requirements and the levels of micronutrients provided in Table 2. These levels include the intrinsic amount of micronutrients in the unfortified milled maize products.

Table 2 — Requirements for micronutrients in fortified milled maize products

SI No	Nutrient	Fortificant compound	Recommended factory level, mg/kg	Regulatory levels, mg/kg	
				Minimum	Maximum
(i)	Vitamin A ¹	Vitamin A (Retinyl) palmitate	1±0.4	0.5	1.4
(ii)	Vitamin B ₁ ¹	Thiamin Mononitrate	6.5± 2.9	3.0	N/A*
(iii)	Vitamin B ₂ ¹	Riboflavin	4±1.8	2	N/A*
(iv)	Niacin ¹	Niacinamide	30±13.4	14.9	N/A*
(v)	Vitamin B ₆ ¹	Pyridoxine	5±2.5	2	N/A*
(vi)	Folate	Folic acid	1.2±0.5	0.6	1.7
(vii)	Vitamin B ₁₂	Vitamin B ₁₂ , WS, 0.1%	0.015±0.007	0.007	N/A*
(viii)	Zinc	Zinc oxide	49±16	33	65
(ix)	Total iron	Total iron	31±10	21	41
(x)	Added Iron	NaFeEDTA	20±10	10	30

*N/A – Not applicable. Setting the maximum level for these nutrients is not necessary because the upper tolerable limit for these nutrients is very high.
¹The addition of these micronutrients is optional in Tanzania.

5.2 Fortificants

Fortificants for use shall be stable compounds conforming to specifications in any of the following documents:

- British Pharmacopoeia (BP),
- Food Chemical Codex (FCC),
- Merck Index (MI),
- United States National Formulary (NF),
- European Pharmacopoeia (Ph Eur),
- United States Pharmacopoeia (USP) or
- FAO/WHO Codex Alimentarius Commission (CAC).

5.3 Premix

The fortificants shall be mixed with diluents or carriers as appropriate to form a premix. Diluents or carriers shall conform to USP, BP, Ph Eur, NF, MI, FAO/WHO, or FCC.

The premix shall be made in such a way that at a given rate of addition to the product, the product shall conform to the requirements in Table 2.

The premix may be formulated to conform to the provisions given in Table 3.

NOTE This premix formulation is designed with minimum nutrient composition and does not take into consideration factory overages in the preparations of the premix.

The premix shall be labelled with the addition rate (that is, the amount of premix to be added to the milled maize product) in grams of premix per metric tonne of maize product and dilution factor.

Where the premix is made in accordance with Table 3, the addition rate shall be 500 g of premix per metric tonne of maize flour.

5.4 Stability of fortificants and premixes

The fortificants and premixes shall have storage stability such that no more than 20 % of its original activity will be lost when stored for 21 days at 45 °C in a well closed container at a level of 2.5 g per kg in milled maize products having the moisture content in the range of 13.5 % - 14.5 %.

The supplier of the premix shall provide the stability data for the fortificants and premixes.

Table 3 – Formulation of premix for addition of micronutrients to milled maize products

Micronutrient	Fortificant compound	Amount of micronutrient to be added to food, mg/kg	Amount of fortificant to be added to food, mg/kg	Amount of fortificant in premix, g/kg	Amount of nutrient in premix, g/kg
Vitamin A	Retinyl palmitate, spray-dried or equivalent, 0.075 % retinol, min.	1.0	13.3	26.7	2
Vitamin B ₁	Thiamin mononitrate, 81 %, min.	4.5	5.6	11.1	9
Vitamin B ₂ (Riboflavin)	Riboflavin, 100 %, min.	3.0	3.0	6.0	6
Vitamin B ₃ (Niacin)	Niacinamide, 99 %, min	25.0	25.3	50.5	50
Vitamin B ₆ (Pyridoxine)	Pyridoxine, 82 %, min.	5.0	6.1	12.2	10
Vitamin B ₉ (Folate)	Folic Acid, 100 %, min.	1.0	1.1	2.2	2
Vitamin B ₁₂	Vitamin B ₁₂ , water-soluble form 0.1 %)	0.015	15.0	30.0	0.03
Iron	NaFeEDTA, 13 % Fe, min.	20	153.8	307.7	37
Zinc	Zinc oxide, 80 %, min.	40	50.0	100.0	80
	Filling material (at least 25 %)		68.3	453.6	
		TOTAL	341.5	1000.0	

6 Food additives

Fortified milled maize (corn) products may contain only the permitted food additives in the Codex Stan 192.

7 Hygiene

7.1 Milled maize (corn) products shall be prepared and handled in accordance with the provisions of appropriate sections of EAS 39.

7.2 The product shall conform to the microbiological limits in Table 4.

Table 4 — Microbiological limits

S/No.	Micro-organism	Maximum limit	Methods of test
i)	Total aerobic count per g	10 ⁵	ISO 4832
ii)	<i>E. Coli</i> per 1 g	Absent	ISO 16649-1
iii)	<i>Salmonella</i> per 25 g	Absent	ISO 6579
iv)	Yeast and Moulds cfu/g	10 ⁴	ISO 7954

S/No.	Micro-organism	Maximum limit	Methods of test
v)	<i>S.aureus</i> per 25 g	10 ²	ISO 6888-1

8 Contaminants

8.1 Heavy metals

Fortified milled maize (corn) products shall conform to those maximum limits for heavy metals established by the Codex Alimentarius Commission as specified in Table 5.

Table 5 — Heavy metals limits

S/No.	Heavy metal	Limits (mg/kg)	Test methods
i)	Arsenic (As)	0.1	ISO 27085
ii)	Lead (Pb)	0.2	ISO 6633
iii)	Cadmium (cd)	0.1	ISO 6561-1 or ISO 6561-2

8.2 Pesticide residues

Fortified milled maize (corn) products shall conform to those maximum residue limits established by the Codex Alimentarius Commission for this commodity.

8.3 Mycotoxins

Fortified milled maize (corn) products shall conform to those maximum mycotoxin limits established by the Codex Alimentarius Commission as specified in Table 6.

Table 6 — Mycotoxins limits

S/No.	Mycotoxins	Limits	Test methods
i)	Total aflatoxins, µg/kg, max	10	ISO 16050.
ii)	Aflatoxins B1, µg/kg, max	5	
iii)	Fumonisin, µg/kg, max	2000	AOAC 2001.04

9 Weights and measures

The fill and the weight of the product shall conform to the weights and measures regulations of the importing partner state.

10 Packaging

Fortified milled maize (corn) products shall be packaged in food grade containers. When the product is packaged in sacks, these must be clean, sturdy and strongly sewn or sealed.

11 Labelling

11.1 General labelling

Labelling shall be done in accordance with EAS 38. At the minimum, the following information shall be displayed:

- a) name of product such as "Whole Maize Meal, Sifted Maize meal , Maize flour or Granulated Maize meal";
- b) the word "Fortified" shall be declared before the name of the product;
- c) name and address of the manufacturer/packer/importer;
- d) brand name/registered trade mark;
- e) batch or code number;
- f) net weight in metric units;
- g) the statement "storage instructions
- h) the statement "Human Food";
- i) country of origin;
- j) date of manufacture;
- k) expiry date;
- l) instructions for disposal of used package.

Each product unit may also be marked with the national food fortification logo, where the industry qualifies to use the mark.

11.2 Nutrition labelling

The amount of micronutrients in the maize flour shall be declared on the label in accordance with CAC/GL 2.

11.3 Nutrition and health claims

Fortified milled maize products may have claims on the importance of the micronutrients in nutrition and health. Such claims when declared shall be consistent with CAC/GL 1 and CAC/GL 23.

12 Methods of sampling

Sampling shall be done in accordance with the ISO 24333.

13 Methods of test

Testing for micronutrients may be conducted using any ECSA¹ or any other internationally recognized test methods

¹ Manual of methods for determining micronutrients in fortified foods. www.a2zproject.org/a2zorg/pdf/Manual_Foods.pdf

Annex A (normative)

Determination of acid insoluble ash

A.1 Reagent

Dilute hydrochloric acid — Approximately 5 N, prepared from concentrated hydrochloric acid.

A.2 Procedure

Weigh accurately about 5 g of the product in a tared, clean and dry porcelain dish. Ignite the material in the dish with the flame of a suitable burner for about one hour. Complete the ignition by keeping in a muffle furnace at 500°C to 600°C until grey ash results. Cool in a desiccator and weigh. Repeat the process of igniting, cooling and weighing at half-hour intervals until the difference between two successive weighings is less than one milligram. Note the lowest mass.

To the ash contained in the porcelain dish add 25 mL of dilute hydrochloric acid, cover with a watch glass and heat on a water bath for 10 minutes. Allow to cool and filter the contents of the dish through Whatman filter paper No. 42 or its equivalent. Wash the filter with water until the washings are free from the acid. Return the filter and the residue to the dish. Keep it in an electric air-oven maintained at 135 ± 2 °C for about 3 hours. Ignite in a muffle furnace at about 550 °C to 600 °C for one hour. Cool the dish in a desiccator and weigh. Repeat the process of igniting in the muffle furnace, cooling and weighing at half-hour intervals until the difference between two successive weighings is less than one milligram. Note the lowest mass.

A.3 Calculation

Acid insoluble ash (on dry basis), per cent by mass

$$= 100 \frac{(m_2 - m)}{(m_1 - m)}$$

where

m_2 = mass in g of the porcelain dish with the acid insoluble ash,

m = mass in g of the empty porcelain dish, and

m_1 = mass in g of the porcelain dish with the dried material taken for the determination of total ash

Bibliography

- [1] Codex Standard 154-1985, Codex Standard for Whole Maize (Corn) Meal
- [2] Codex Standard 155-1985 Codex Standard for Degermed Maize (Corn) Meal and Maize (Corn) Grits
- [3] ECSA-HC Guidelines of fortification levels for staples

